



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## MECHANICS.

When this issue was made up solutions had been received for numbers 292, 293, 295-6-7-8-9. Please give attention to 272, 277-8-9, 286-7, 290-1, 294, 300.

**301. Proposed by CLIFFORD N. MILLS, Brookings, S. D.**

A wire is hanging from two points in the same horizontal plane. If the difference between the length of the wire and the actual distance between the supports is very small, show that

$$s = x \left( 1 + \frac{x^2}{6c^2} \right),$$

where  $s$  is one half the length of the wire,  $c$  is the horizontal tension at the lowest point divided by  $w$  the load per unit of horizontal distance, and  $x$  is the distance of lowest point of the curve to the point of support.

## NUMBER THEORY.

When this issue was made up solutions had been received for numbers 212-13, 215-16, 218, 220, 223. Please give attention to 191-2, 196, 198, 201-2, 205, 208-9-10-11, 214, 217, 219, 221-2.

**224. Proposed by PATRICK WALSH, New Orleans, Louisiana.**

Find the sides, in rational numbers, of a right angled triangle whose area is  $5\frac{1}{2}$ .

**225. Proposed by W. DE W. CAIRNS, Oberlin College.**

*L'Intermédiaire* for June, 1914, contains the following problem:

"If we write the terms of the arithmetic series 1, 5, 9, 13, 17, 21, 25, 29, 33, ... as follows:

$$\begin{array}{ccccccc} 1 & & & & & & \\ 5 & 9 & 13 & & & & \\ 17 & 21 & 25 & 29 & 33 & & \\ 37 & 41 & 45 & 49 & 53 & 57 & 61 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{array},$$

it is seen that the sum of the terms of each line is a cube, and that these are the cubes of the successive "odd integers. How is this shown?"

It is here proposed not only to prove this, but to generalize the theorem as suggested, using, however, the simpler (and better known) case which includes all of the successive integers:

$$\begin{array}{cccc} 1 & & & \\ 3 & 5 & & \\ 7 & 9 & 11 & \\ 13 & 15 & 17 & 19 \\ \cdot & \cdot & \cdot & \cdot \end{array}$$

## SOLUTIONS OF PROBLEMS.

## ALGEBRA.

**412. Proposed by H. L. SLOBIN, University of Minnesota.**

Form the algebraic equation whose roots are:

$$a_1 = \cos \frac{\pi}{9}, \quad a_2 = -\cos \frac{2\pi}{9}, \quad a_3 = -\cos \frac{4\pi}{9}.$$

SOLUTION BY CLIFFORD N. MILLS, Brookings, South Dakota.

From the suggestions given by the proposer,

$$\cos r\pi = \frac{(-1)^r + (-1)^{-r}}{2}.$$